Data Science Toolbox Question Sheet

02.1 Regression

Daniel Lawson

Block 2

Short questions

1. In the univariate case with K = 1, the correlation between two samples from the same distribution is written

$$\operatorname{Corr}(X_1, X_2) = \frac{\mathbb{E}[(X_1 - \mu)(X_2 - \mu)]}{\sigma_X^2}.$$

Define multivariate correlation in matrix notation.

- 2. X and Y are found to be correlated, conditional on some additional variables Z. Under what circumstances does this imply that X causes Y?
- 3. Consider the linear model $\mathbf{y} = \mathbf{x}\beta + \epsilon$ where \mathbf{x} is of dimension S. Using $(a + bx)^T = (a^T + x^T b^T)$ and $a = a^T$ when a is a 1 by 1 matrix (or otherwise), demonstrate that:

$$MSE(\beta) = \frac{1}{n} (\mathbf{y}^T \mathbf{y} - 2\beta^T x^T \mathbf{y} + \beta^T \mathbf{x}^T \mathbf{x}\beta).$$

- 4. Recall that $\hat{H} = X((X^TX)^{-1})X^T$. Show that $\mathbb{I} \hat{H}$ is symmetric, by computing \hat{H}^T or otherwise.
- 5. Compute $\hat{H}\hat{H}$ to show that it is Idempotent.
- 6. Discuss the value of unbiasedness of the estimator for $\hat{\beta}$, i.e. that $\mathbb{E}(\hat{\beta}) = \beta$.
- 7. Prove that $\mathbb{E}(\mathbf{y}) = \mathbf{x}\beta$.
- 8. Compute $Var(\mathbf{y})$ and discuss the circumstances in which it simplifies.